

DEFORMATIONS OF SPECIAL LEGENDRIAN SUBMANIFOLDS WITH BOUNDARY

GUANGCUN LU AND XIAOMIN CHEN

ABSTRACT. In this article, for a compact special Legendrian submanifold with boundary of contact Calabi-Yau manifolds we study the deformation of it with boundary confined in an appropriately chosen contact submanifold of codimension two which we also a scaffold (Definition 2.3) by analogy with [A.Butsher, Deformations of minimal Lagrangian submanifolds with boundary, Proc. Amer. Math. Soc. 131(2002) 1953-1964]. Our first result shows that it cannot be deformed, and the second claims that deformations of such a special Legendrian submanifold forms a one-dimensional smooth manifold under suitably weaker boundary confinement conditions. They may be viewed as supplements of the boundless case considered by Tomassini and Vezzoni [Contact Calabi-Yau manifolds and special Legendrian submanifolds, Osaka J. Math., 45(2008), 127-147].

1. INTRODUCTION AND MAIN RESULTS

The calibrated geometry was invented by Harvey and Lawson in their seminal paper [5]. A class of important calibrated submanifolds is special Lagrangian submanifolds in Calabi-Yau manifolds. Let (M, J, ω, Ω) be a real $2n$ -dimensional Calabi-Yau manifold. A special Lagrangian submanifold of it is a submanifold L with $\omega|_L = 0$ and $\text{Im}(\Omega)|_L = 0$. In 1996 McLean [10] developed the deformation theory of special Lagrangian submanifolds (and other special calibrated submanifolds) and showed:

McLean Theorem ([10]). *A normal vector field V to a compact boundaryless special Lagrangian submanifold L in (M, J, ω, Ω) is the deformation vector field to a normal deformation through special Lagrangian submanifolds if and only if the corresponding 1-form $(JV)^\flat$ on L is harmonic. There are no obstructions to extending a first order deformation to an actual deformation and the tangent space*

Date: June 25, 2011.

2000 Mathematics Subject Classification. Primary 53C10, 53D10, 53C25, 53C38.

Key words and phrases. Contact Calabi-Yau manifolds, special Legendrian submanifolds with boundary, scaffold.

Partially supported by the NNSF 10671017 and 10971014 of China, and PCSIRT and Research Fund for the Doctoral Program Higher Education of China (Grant No. 200800270003).

to such deformations can be identified through the cohomology class of the harmonic form with $H^1(L; \mathbb{R})$.

Since then the theory is generalized to various situations. See [6, 7, 13] and references therein. For example, S. Salur [14] generalized McLean theorem to symplectic manifolds. We here only list those closely related to ours. The first one is the case of compact special Lagrangian submanifolds with nonempty boundary considered by Butsher [1]. He called a submanifold L in the Calabi-Yau manifold (M, J, ω, Ω) **minimal Lagrangian** if $\omega|_L = 0$ and $\text{Im}(e^{i\theta}\Omega)|_L = 0$ for some $\theta \in \mathbb{R}$. If L is a Lagrangian submanifold of (M, ω) with nonempty boundary ∂L and $N \in \Gamma(T_{\partial L}L)$ is the inward unit normal vector field of ∂L in L , he defined a **scaffold** for L to be a submanifold W of M such that $\partial L \subset W$, the bundle $(TW)^\omega$ is trivial, and that N is a smooth section of the bundle $(T_{\partial L}W)^\omega$.

Butsher Theorem ([1]). *Let L be a special Lagrangian submanifold of a compact Calabi-Yau manifold M with non-empty boundary ∂L and let W be a symplectic, codimension two scaffold for L . Then the space of minimal Lagrangian submanifolds sufficiently near L (in a suitable $C^{1,\beta}$ sense) but with boundary on W is finite dimensional and is parametrized over the harmonic 1-forms of L satisfying Neumann boundary conditions.*

The work inspired Kovalev and Lotay [8] to study the analogous deformation problem of a compact coassociative 4-fold with boundary inside a fixed coassociative in a 7-manifold with closed G_2 -structures. Recently Gayet and Witt [3] also investigated the deformation of a compact associative submanifold with boundary in a coassociative submanifold in a topological G_2 -manifold.

As a natural generalization of the Calabi-Yau manifolds in the context of contact geometry Tomassini and Vezzoni [17, Def.3.1] introduced the notion of a contact Calabi-Yau manifold, cf. Definition 2.1. Let (M, η, J, ϵ) be a $(2n+1)$ -dimensional contact Calabi-Yau manifold, and $j : L \hookrightarrow M$ be a compact special Legendrian submanifold without boundary (cf. Definition 2.2). Two special Legendrian submanifolds $j_0 : L \hookrightarrow M$ and $j_1 : L \hookrightarrow M$ are called **deformation equivalent** if there exists a smooth map $F : L \times [0, 1] \rightarrow M$ such that

- $F(\cdot, t) : L \times \{t\} \rightarrow M$ is a special Legendrian embedding for any $t \in [0, 1]$;
- $F(\cdot, 0) = j_0$, $F(\cdot, 1) = j_1$.

(cf. [17, Def.4.4]). If there exists a diffeomorphism $\phi \in \text{Diff}(L)$ such that $j_1 = j_0 \circ \phi$ we say j_0 and j_1 to be **equivalent**. This yields an equivalent relation $\sim$ among all embeddings from L to M . Let $\widetilde{\mathfrak{M}}(L)$ be the set of special Legendrian submanifolds of (M, α, J, ϵ) which are deformation equivalent to $j : L \hookrightarrow M$. Call $\mathfrak{M}(L) := \widetilde{\mathfrak{M}}(L) / \sim$ the **Moduli space of special Legendrian submanifolds** which are deformation equivalent to $j : L \hookrightarrow M$. Tomassini and Vezzoni [17,

Theorem 4.5] proved:

Tomassini-Vezzoni Theorem([17, Theorem 4.5]). *Let (M, η, J, ϵ) be a contact Calabi-Yau manifold of dimension $(2n + 1)$, and $L \subset M$ be a compact special Legendrian submanifold without boundary. Then the moduli space $\mathfrak{M}(L)$ is a smooth one-dimensional manifold.*

Motivated by the above works, we study in this paper the local deformations of compact special Legendrian submanifolds with (nonempty) boundary. Different from the case $\partial L = \emptyset$ considered by Tomassini and Vezzoni [17], it is showed in Remark 5.1 that the moduli space $\mathfrak{M}(L)$ is infinite dimensional.

In order to get intersecting results it is necessary to add some boundary conditions. Inspired by [1, Def. 1] we introduce a notion of **scaffold** for L in Definition 2.3, which is a suitable contact submanifold W of codimension two. Two special Legendrian submanifolds $j_0 : L \hookrightarrow M$ and $j_1 : L \hookrightarrow M$ with $j_0(\partial L) \subset W$ and $j_1(\partial L) \subset W$ are called **deformation equivalent** if there exists a smooth map $F : L \times [0, 1] \rightarrow M$ such that

- $F(\cdot, t) : L \times \{t\} \rightarrow M$ is a special Legendrian embedding with $F(\partial L, t) \subset W$ for any $t \in [0, 1]$;
- $F(\cdot, 0) = j_0, F(\cdot, 1) = j_1$.

The **Moduli space of special Legendrian submanifolds** which are deformation equivalent to $j : L \hookrightarrow M$ with $j(\partial L) \subset W$ is defined as

$$\begin{aligned} \mathfrak{M}(L, W) := \{ & \text{special Legendrian submanifolds of } (M, \alpha, J, \epsilon) \\ & \text{which are deformation equivalent to } j : L \hookrightarrow M \\ & \text{with } j(\partial L) \subset W \text{ and are near } j \} / \sim . \end{aligned}$$

Our first result is

Theorem 1.1. *Let (M, J, α, ϵ) be a contact Calabi-Yau manifold, and L be a compact special Legendrian submanifold with nonempty boundary ∂L inside a scaffold W of codimension two. Then L cannot be deformed as a special Legendrian submanifold with boundary confined in W . In other words $\mathfrak{M}(L, W)$ only consists of the class of j .*

This is in contrast with the case of compact boundless special Legendrian submanifolds considered in Tomassini-Vezzoni Theorem. Such a local rigidity is similar to the case of a compact simply connected boundless special Lagrangian submanifold in McLean Theorem, and a stable result of minimal submanifolds proved by Simons [16].

Now we turn to consider weaker boundary conditions. Let (M, α, J, ϵ) be a $(2n + 1)$ -dimensional contact Calabi-Yau manifold, and $L \subset M$ be a compact

special Legendrian submanifold with (non-empty) boundary. A normal vector field V to L is called **boundary α -constant** if $\alpha(V)|_{\partial L}$ is constant. The following result, which is stated in a similar way to McLean Theorem above, is similar to that of Tomassini and Vezzoni [17].

Theorem 1.2. *Let (M, α, J, ϵ) be a $(2n+1)$ -dimensional contact Calabi-Yau manifold, and $L \subset M$ be a compact special Legendrian submanifold with (non-empty) boundary. A boundary α -constant normal vector field V to L is the deformation vector field to a normal deformation through special Legendrian submanifolds if and only if $\alpha(V)$ is constant. Moreover the tangent space to such deformations can be identified with $\mathbb{R}$.*

The key points in the proof of Theorems 1.1 and 1.2 are to find a suitable definition of scaffold for a special Legendrian submanifold with boundary and to prove a corresponding result with Lemma 5 of [1], Lemma 3.1. For the former we propose and study it in Section 2. The proof of the latter will be given in Section 3 and is more troublesome because we need to use not only contact neighborhood theorem but also symplectic neighborhood theorem. In Sections 4 and 5, we complete the proofs of Theorems 1.1 and 1.2 respectively.

2. PRELIMINARIES

2.1. Contact Calabi-Yau manifolds and special Legendrian submanifolds.

Let (M, α) be a contact manifold with contact distribution $\xi = \ker \alpha$ and Reeb vector field R_α . Then $\kappa := d\alpha/2$ restricts a symplectic vector bundle structure on $\xi \rightarrow M$, $\kappa|_\xi$, and every compatible complex structure $J \in \mathcal{J}(\xi, \kappa|_\xi)$ gives a Riemannian metric g_J on the bundle $\xi \rightarrow M$, $g_J(u, v) = \kappa(u, Jv)$ for $u, v \in \xi$. By setting $J(R_\alpha) = 0$ we can extend J to an endomorphism of TM , also denoted by J without special statements. Clearly

$$J^2 = -\mathbf{I} + \alpha \otimes R_\alpha, \quad \text{and} \quad g := g_J + \alpha \otimes \alpha \quad (2.1)$$

is a Riemannian metric g on M , where $\mathbf{I}$ is the identity endomorphism on TM . Define a Nijenhuis tensor of J by

$$N_J(X, Y) = [JX, JY] - J[X, JY] - J[JX, Y] + J^2[X, Y]$$

for all $X, Y \in TM$. If $N_J = -d\alpha \otimes R_\alpha$ then the pair (α, J) is a **Sasakian structure** on M , and the triple (M, α, J) is called a **Sasakian manifold**. On such a manifold it holds that $d\Lambda_B^r(M) \subset \Lambda_B^r(M)$ and $J(\Lambda_B^r(M)) = \Lambda_B^r(M)$, where $\Lambda_B^r(M)$ is the set of all differential r -form γ on M with $\iota_{R_\alpha}\gamma = 0$ and $\mathcal{L}_{R_\alpha}\gamma = 0$. So we have a split

$$\Lambda_B^r(M) \otimes \mathbb{C} = \oplus_{p+q=r} \Lambda_J^{p,q}(\xi)$$

and $\kappa = \frac{1}{2}d\alpha \in \Lambda_J^{1,1}(\xi)$.

Definition 2.1 ([17, Def.2.1]). A **contact Calabi-Yau manifold** is a quadruple (M, α, J, ϵ) consisting of a $(2n+1)$ -dimensional Sasakian manifold (M, α, J) and a nowhere vanishing basic form $\epsilon \in \Lambda_J^{n,0}(\xi)$ such that

$$\epsilon \wedge \bar{\epsilon} = c_n \frac{\kappa^n}{n!} \quad \text{and} \quad d\epsilon = 0,$$

where $c_n = (-1)^{\frac{n(n+1)}{2}} (2i)^n$ and $\kappa = \frac{1}{2}d\alpha$.

Definition 2.2 ([17, Def.4.2]). Let $(M^{2n+1}, \alpha, J, \epsilon)$ be a contact Calabi-Yau manifold. An embedding $p : L \rightarrow M$ is called a **special Legendrian submanifold** if $\dim L = n$, $p^*\alpha = 0$ and $p^*\text{Im}\epsilon = 0$.

Clearly, $p^*\epsilon = p^*(\text{Re}\epsilon)$ is a volume form on L . Thus every special Legendrian submanifold has a natural orientation. By [10, p.722] or [2, Prop.2.6] we have

$$p^*(\iota_Y \text{Im}\epsilon) = -\star(p^*(\iota_Y \kappa)) = -\frac{1}{2}\star(p^*(\iota_Y d\alpha)) \quad (2.2)$$

for any section $Y : L \rightarrow p^*\xi$, where the star operator $\star$ is computed with respect to $p^*(g_J) = p^*(\kappa \circ (\text{id} \times J))$ and the volume form $\text{Vol}(L) := p^*\epsilon = p^*(\text{Re}\epsilon)$.

For any n -dimensional manifold N , the cotangent bundle T^*N has a canonical 1-form λ_{can} . The 1-jet bundle $J^1N = \mathbb{R} \times T^*N$ is a contact manifold with contact form $\alpha = \pi_1^*(dt) - \pi_2^*(\lambda_{\text{can}})$ and Reeb vector field $\partial/\partial t$, where $t \in \mathbb{R}$ is the real parameter and π_i is the projection from $\mathbb{R} \times T^*N$ onto the i -th factor, $i = 1, 2$. (See [9, Example 3.44]).

2.2. Boundary conditions. Corresponding to [1, Def. 1] we introduce:

Definition 2.3. Let L be a submanifold of the contact manifold $(M, \xi = \ker \alpha)$ with boundary ∂L and let $N \in \Gamma(T_{\partial L}L)$ be the inward unit normal field of ∂L in L . A contact submanifold (W, ξ') of (M, ξ) is called a **scaffold** for L if (i) $\partial L \subset W$, (ii) $N \in \Gamma(\xi'^{\perp}|_{\partial L})$, and (iii) the bundle $\xi'^{\perp}$ is trivial, where $\xi'^{\perp}$ is the symplectically orthogonal complement of ξ' in $(\xi|_W, \kappa|_{\xi|_W})$.

Given a contact manifold (M, α) let J and g be as in (2.1). Denote by $\perp_g$ orthogonal with respect to the compatible metric g on M . If (W, ξ') is a contact submanifold of $(M, \xi = \ker \alpha)$, that is, $T_x W \cap \xi_x = \xi'_x$ for all $x \in W$, the following claim the condition (iii) of Definition 2.3 is equivalent to one that $(TW)^{\perp_g}$ is trivial.

Claim 2.4. $(TW)^{\perp_g} = (J\xi')^{\perp} = J(\xi'^{\perp})$.

Proof. For $x \in W$, since $\xi_x'^\perp \subset \xi_x$ and J_x restricts to a complex structure ξ_x we have

$$\begin{aligned}
\xi_x'^\perp &= \{v \in \xi_x \mid \kappa(v, u) = 0 \ \forall u \in \xi_x'\} \\
&= \{v \in \xi_x \mid \kappa(Jv, Ju) = 0 \ \forall u \in \xi_x'\} \\
&= \{v \in \xi_x \mid g_J(Jv, u) = 0 \ \forall u \in \xi_x'\} \\
&= \{v \in \xi_x \mid g(Jv, bR_\alpha + u) = 0 \ \forall bR_\alpha + u \in R_\alpha \mathbb{R} + \xi_x'\} \\
&= \{v \in \xi_x \mid g(Jv, Y) = 0 \ \forall Y \in T_x W\}.
\end{aligned}$$

This implies $J\xi'^\perp = (TW)^\perp_g$ or $\xi'^\perp = J(TW)^\perp_g$. Moreover, both $J\xi'^\perp$ and $\xi'^\perp$ are contained in $\xi|_W$, and ξ is J -invariant. It is easy to check that $J\xi'^\perp = (J\xi')^\perp$. $\square$

Proposition 2.5. *Let L be a Legendrian submanifold of the contact manifold $(M, \xi = \ker \alpha)$ with (nonempty) boundary ∂L and W be a scaffold for L . Then ∂L is a Legendrian submanifold of (W, ξ') .*

Proof. Since L is the Legendrian submanifold of (M, ξ) , $TL \subset \xi|_L$. Moreover the definition of the scaffold implies that $T\partial L \subset T_{\partial L}W$ and thus $T\partial L \subset T_{\partial L}W \cap \xi|_{\partial L} = \xi'|_{\partial L}$. This shows that the boundary ∂L is a Legendrian submanifold of (W, ξ') . $\square$

Under the assumptions of Proposition 2.5, let $f_t : L \rightarrow M$ be a deformation of L satisfying $f_t(\partial L) \subset W$ for all t , and let $V = \frac{d}{dt}f_t|_{t=0}$ be the corresponding deformation vector field. Clearly, $V(x) \in T_x W$ for any $x \in \partial L$. Since L is a Legendrian submanifold, we have $TL \subset \xi|_L$. Note that $N(x) \in T_x L$ for any $x \in \partial L$. Then the condition (ii) of Definition 2.3 implies that $N(x) \in \xi_x^\perp$, and so $N(x) \in T_x L \cap \xi_x'^\perp$ and

$$J_x N(x) \in J_x(T_x L \cap \xi_x'^\perp) \subset J_x \xi_x'^\perp \subset J_x \xi_x = \xi_x.$$

Since W is a contact submanifold, we may write $V(x) = Y + aR_\alpha(x)$, where $Y \in \xi'_x$. By Claim 2.4, $J_x N(x) \in J_x \xi_x'^\perp = (T_x W)^\perp_g$ and thus

$$0 = g(J_x N(x), V(x)) = g_J(J_x N(x), Y) = \kappa(J_x N(x), J_x Y) = \kappa(N(x), Y).$$

Note that $Y = V(x) - \alpha(V(x))R_\alpha(x)$ and that $\iota_{R_\alpha} d\alpha = 0$. We get

Claim 2.6. *If $f_t : L \rightarrow M$ be a deformation of L satisfying $f_t(\partial L) \subset W$ for all t , then the corresponding deformation vector field V satisfies*

Neumann boundary condition: $d\alpha(N(x), V(x)) = 0 \ \forall x \in \partial L$.

3. CONSTRUCTING A NEW METRIC

In the study of the deformation of the special Legendrian submanifold L without boundary by Tomassini and Vezzoni [17], the deformations of L are parameterized by sections of the normal bundle $N(L)$ using the exponent map

$\exp(V) : L \rightarrow M$. However, in our case, since W is generally not totally geodesic submanifold, it cannot be assured that the image of ∂L under $\exp(V)$ sits in W . In order to fix out the problem we shall follow the ideas in [1] to construct a new metric $\hat{g}$ such that the image of ∂L under the corresponding exponent map is contained in W , that is, such that W is totally geodesic. The following is an analogue of [1, Lemma 5].

Lemma 3.1. *Let L be a compact Legendrian submanifold of the contact manifold (M, J, α) with (nonempty) boundary ∂L and W be a scaffold for it of codimension two. Then there is a neighborhood $\mathcal{U} = \mathcal{U}(\partial L, M)$ of ∂L in M and a contact embedding $\phi : \mathcal{U} \rightarrow \mathbb{R} \times T^*(\partial L) \times \mathbb{R}^2$ such that the following conclusions hold:*

- (i) $\phi(W \cap \mathcal{U}) \subset \mathbb{R} \times T^*(\partial L) \times \{(0, 0)\}$,
- (ii) $\phi(\partial L) = \{0\} \times \partial L \times \{0, 0\}$,
- (iii) $(t, x, v, s_1, s_2) \in \phi(\mathcal{U}) \implies (t, x, v, 0, 0) \in \phi(\mathcal{U})$,
- (iv) *For any nowhere zero smooth section $V : W \rightarrow \xi'^\perp|_W$, ϕ can be required to satisfy $\phi_*(V(p)) = \frac{\partial}{\partial s_1} \Big|_{\phi(p)}$ for any $p \in \partial L$, where (s_1, s_2) the coordinate functions of $\mathbb{R}^2$.*

Note that the conclusion (iv) is slightly weaker than the corresponding one of [1, Lemma 5(4)]. It is sufficient for us to construct a suitable metric in Proposition 3.2. Even so our proof uses not only contact neighborhood theorem but also symplectic neighborhood theorem in contrast with the proof of [1, Lemma 5(4)]. It is a key of our proof.

Proof of Lemma 3.1. Since ∂L is a compact Legendrian submanifold of $\mathbb{R} \times T^*(\partial L)$ without boundary, from the Neighborhood Theorem for Legendrian (cf. Corollary 2.5.9 in [4]) it follows that there exists a contactomorphism ϕ_0 from a neighborhood $\mathcal{U}_0(\partial L, W)$ of ∂L in W to one $\mathcal{V}_0(0_{\partial L})$ of the zero section of $T^*(\partial L)$ in $\mathbb{R} \times T^*(\partial L)$ such that

$$\phi_0(x) = (0, x) \quad \forall x \in \partial L. \quad (3.1)$$

Fix a Riemannian metric on the bundle $T^*(\partial L)$, and then take a sufficiently small $\epsilon > 0$ such that

$$M'_1 := \{(t, x, v) : |t| \leq \epsilon, v \in T_x^*(\partial L) \text{ with } |v| \leq \epsilon\} \subset \mathcal{V}_0(0_{\partial L}). \quad (3.2)$$

We get another neighborhood of ∂L in W ,

$$M'_0 := \phi_0^{-1}(M'_1) \subset \mathcal{U}_0(\partial L, W) \subset W. \quad (3.3)$$

Then $\phi_0 : M'_0 \rightarrow M'_1$ is a contactomorphism. Obverse that M'_0 and M'_1 are compact contact submanifolds of W and $T^*(\partial L) \times \mathbb{R}$ with boundary and of codimension zero, respectively.

Let λ_{can} denote the canonical 1-form on $T^*\partial L$. Recall that the contact form and Reeb vector field on $J^1\partial L = \mathbb{R} \times T^*(\partial L)$ are

$$\tilde{\beta} = dt - \lambda_{\text{can}} \quad \text{and} \quad R_{\tilde{\beta}} = \partial/\partial t. \quad (3.4)$$

Assume that s_1, s_2 are the coordinate functions of $\mathbb{R}^2$. We have a contact form on $J^1\partial L \times \mathbb{R}^2 = \mathbb{R} \times T^*(\partial L) \times \mathbb{R}^2$,

$$\beta = \tilde{\beta} - s_1 ds_2 = dt - \lambda_{\text{can}} - s_1 ds_2, \quad (3.5)$$

whose Reeb vector field is given by $R_\beta = \partial/\partial t$. Denote by $(\ker(\tilde{\beta}))^\perp$ the symplectically orthogonal complement of $\ker(\tilde{\beta})$ in $\ker(\beta)$ (with respect to $d\beta$). It is easily checked that it is equal to the trivial bundle

$$\text{Span}\left(\left\{\frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_2}\right\}\right) \rightarrow J^1\partial L \times \mathbb{R}^2.$$

Define $M_0 := M$, $M_1 := \mathbb{R} \times T^*(\partial L) \times \mathbb{R}^2$, and M'_0 and M'_1 as above. (Identify $M'_1 \equiv M'_1 \times \{(0, 0)\} \subset J^1L \times \mathbb{R}^2$). Since $\xi'^\perp$ is trivial we can pick two vector fields V_1, V_2 such that V_1, V_2 form a basis of $\xi'^\perp$ and satisfy $d\alpha(V_1, V_2) = 0$. There exists an obvious symplectic vector bundle isomorphism

$$\xi'^\perp|_{M'_0} \rightarrow \text{Span}\left(\left\{\frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_2}\right\}\right) \Big|_{M'_1}$$

given by

$$\Phi(V_1(x)) = \frac{\partial}{\partial s_1} \Big|_{(\phi_0(x), 0, 0)} \quad \text{and} \quad \Phi(V_2(x)) = \frac{\partial}{\partial s_2} \Big|_{(\phi_0(x), 0, 0)}$$

for any $x \in M'_0$. By Theorem 2.5.15 of [4], we may extend ϕ_0 into a contactomorphism ϕ_1 from a neighborhood $\mathcal{U}(M'_0)$ of M'_0 in $M_0 = M$ to that $\mathcal{U}(M'_1)$ of $M'_1 \equiv M'_1 \times \{(0, 0)\}$ in M_1 such that $T\phi_1|_{\xi'^\perp|_{M'_0}}$ and Φ are bundle homotopic (as symplectic bundle isomorphisms) up to a conformality. (**Note:** From the proof of [4, Th.2.5.15] it is not hard to see that the theorem still holds if compact contact submanifold M'_i have boundary and $M'_i \subset \text{Int}(M_i)$.)

Actually, we may assume that $\mathcal{U}(M'_1)$ has the following form:

$$\begin{aligned} \mathcal{U}(M'_1) &= \{(t, x, v) : |t| < \varepsilon, v \in T_x^*(\partial L) \text{ with } |v| < \varepsilon'\} \\ &\quad \times \{(s_1, s_2) \in \mathbb{R}^2 : |s_1|, |s_2| < \delta\}, \end{aligned} \quad (3.6)$$

where $0 < \varepsilon' < \varepsilon$ and $\delta > 0$, and

$$\mathcal{U}(M'_0) := \phi_1^{-1}(\mathcal{U}(M'_1)).$$

By suitably shrinking $\mathcal{U}(M'_0)$ and $\mathcal{U}(M'_1)$ if necessary, we can require

$$W_0 := W \cap \mathcal{U}(M'_0) \subset \mathcal{U}_0(\partial L, W), \quad (3.7)$$

$$\phi_1(W_0) \subset \mathbb{R} \times T^*\partial L \times \{(0, 0)\}, \quad (3.8)$$

$$(t, x, v, s_1, s_2) \in \mathcal{U}(M'_1) \implies \phi_1^{-1}(t, x, v, 0, 0) \in W_0.$$

Clearly, $\mathcal{U}(M'_0)$ and ϕ_1 satisfy the conditions (i)-(iii) in Lemma 3.1.

For (iv) we need to modify ϕ_1 and $\mathcal{U}(M'_0)$. Since ϕ_1 is a contactomorphism,

$$\phi_{1*}(\zeta'^\perp|_{W_0}) \subset \text{Span}\left(\left\{\frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_2}\right\}\right) \Big|_{\phi_1(W_0)}.$$

It follows that there exist smooth real functions $f_1, f_2 : W_0 \rightarrow \mathbb{R}$ such that

$$\phi_{1*}(V(x)) = f_1(x) \frac{\partial}{\partial s_1} \Big|_{\phi(x)} + f_2(x) \frac{\partial}{\partial s_2} \Big|_{\phi(x)} \quad \text{and} \quad |f_1(x)| + |f_2(x)| \neq 0$$

for any $x \in W_0$.

Take $\epsilon > 0$ sufficiently small so that

$$R_\epsilon := \{(t, x, v, 0, 0) \in \mathbb{R} \times T^*\partial L \times \mathbb{R}^2 : |t| \leq \epsilon, |v| \leq \epsilon\} \subset \phi_1(W_0).$$

Consider the compact symplectic submanifold of $(T^*\partial L \times \mathbb{R}^2, -d\lambda_{\text{can}} - ds_1 \wedge ds_2)$,

$$S_\epsilon := \{(x, v, 0, 0) \in T^*\partial L \times \mathbb{R}^2 : |v| \leq \epsilon\}. \quad (3.9)$$

Its symplectic normal bundle is

$$\text{Span}\left(\left\{\frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_2}\right\}\right) \Big|_{S_\epsilon},$$

and $\phi_{1*}(V)$ restricts to a nowhere zero smooth section

$$p \mapsto f_1 \circ \phi^{-1}(p) \frac{\partial}{\partial s_1} \Big|_p + f_2 \circ \phi^{-1}(p) \frac{\partial}{\partial s_2} \Big|_p. \quad (3.10)$$

Obverse that there exists an obvious symplectic vector bundle isomorphism

$$\Psi : \text{Span}\left(\left\{\frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_2}\right\}\right) \Big|_{S_\epsilon} \rightarrow \text{Span}\left(\left\{\frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_2}\right\}\right) \Big|_{S_\epsilon}$$

which sends the section in (3.10) to one

$$p \mapsto \frac{\partial}{\partial s_1} \Big|_p.$$

Hence the symplectic neighborhood theorem¹ (cf. [9, Th.3.30]) yields a symplectomorphism between neighborhoods of S_ϵ in $(T^*\partial L \times \mathbb{R}^2, -d\lambda_{\text{can}} - ds_1 \wedge ds_2)$,

$$\varphi : \mathcal{N}_0(S_\epsilon) \rightarrow \mathcal{N}_1(S_\epsilon)$$

¹From the proof of [9, Th.3.30] it is not hard to see that the theorem still holds if compact symplectic submanifold Q_j have boundary and $Q_j \subset \text{Int}(M_j)$.

such that

$$\varphi(p) = p \quad \text{and} \quad d\varphi(p) = \Psi_p \quad (3.11)$$

for any $p \in S_\epsilon$. In particular, we have

$$d\varphi(p)\left(\phi_{1*}(V)|_p\right) = \frac{\partial}{\partial s_1} \Big|_p \quad \forall p \in S_\epsilon. \quad (3.12)$$

Since (3.5) implies

$$\ker(\beta) \Big|_{(t,x,v,s_1,s_2)} = T_{(x,v)}T^*\partial L \times \text{Span}\left(\left\{\frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_2}\right\}\right) \Big|_{(s_1,s_2)},$$

the map

$$\phi_2 : \mathbb{R} \times \mathcal{N}_0(S_\epsilon) \rightarrow \mathbb{R} \times \mathcal{N}_1(S_\epsilon), \quad (t, p) \mapsto (t, \varphi(p)) \quad (3.13)$$

must be a contactomorphism with respect to the induced contact structure from $(\mathbb{R} \times T^*(\partial L) \times \mathbb{R}^2, \beta)$.

Take a neighborhood $\mathcal{U}$ of ∂L in M such that

$$\begin{aligned} \mathcal{U} &\subset \mathcal{U}(M'_0) \quad \text{and} \quad \phi_1(\mathcal{U}) \subset \mathbb{R} \times \mathcal{N}_0(S_\epsilon), \\ (t, x, v, s_1, s_2) &\in \phi_2(\phi_1(\mathcal{U})) \implies (t, x, v, 0, 0) \in \phi_2(\phi_1(\mathcal{U})). \end{aligned}$$

Then the composition $\phi := \phi_2 \circ (\phi_1|_{\mathcal{U}})$ is a contact embedding from $\mathcal{U}$ into $(\mathbb{R} \times T^*(\partial L) \times \mathbb{R}^2, \beta)$ such that the condition (iii) is satisfied. By (3.8) and (3.11) it is easy to see that (i) is satisfied for ϕ and $\mathcal{U}$, i.e.

$$\phi(W \cap \mathcal{U}) \subset \mathbb{R} \times T^*\partial L \times \{(0, 0)\}.$$

From (3.1) it follows that $\phi(\partial L) = \{0\} \times \partial L \times \{0, 0\}$. That is, (i) holds. Finally, (3.12) implies that ϕ satisfies the condition (iv), i.e.

$$d\phi(p)(V(p)) = \frac{\partial}{\partial s_1} \Big|_{\phi(p)} \quad \forall p \in \partial L.$$

□

As in [1], with Lemma 3.1 we may construct the desired metric $\hat{g}$ as follows.

Step 1. Recall that N the unit normal vector field of ∂L in L . Let $\mathcal{U}$ and ϕ be as in the Lemma 3.1 with $\phi_*(N(p)) = \frac{\partial}{\partial s_1} \Big|_{\phi(p)}$ for any $p \in \partial L$. By Lemma 3.1(iii) we may define a metric g' on $\phi(\mathcal{U})$ as follows:

$$g'(t, x, v, s_1, s_2) := (\phi^{-1})^*(g|_W(\phi^{-1}(t, x, v, 0, 0))) + ds_1 \otimes ds_1 + ds_2 \otimes ds_2$$

for every $(t, x, v, s_1, s_2) \in \phi(\mathcal{U})$.

Step 2. Consider the metric $g_1 := \phi^*g'$ on $\mathcal{U}$. Take a neighborhood $\mathcal{V}$ of ∂L in M such that the closure of $\mathcal{V}$ is contained in $\mathcal{U}$. Let $\rho : M \rightarrow \mathbb{R}$ be a smooth function such that $\rho = 1$ on a neighborhood $\mathcal{V}$, and $\rho = 0$ outside $\mathcal{U}$. We then define the metric $\hat{g}$ by

$$\hat{g} := \rho g' + (1 - \rho)g.$$

The following two propositions correspond to Propositions 6 and 7 in [1], respectively.

Proposition 3.2. *For the neighborhood $\mathcal{V}$ of ∂L in Step 2, $W \cap \mathcal{V}$ is totally geodesic with respect to the metric $\hat{g}$.*

Proof. For any $p \in W \cap \mathcal{V}$, Lemma 3.1 gives a local contact coordinate system around it,

$$\mathcal{O}(p) \rightarrow \mathbb{R} \times \mathbb{R}^{2n-2} \times \mathbb{R}^2, \quad q \mapsto (t(q), z_1(q), \dots, z_{2n-2}(q), s_1(q), s_2(q))$$

such that

- for some smooth function $h : \mathcal{O}(p) \rightarrow \mathbb{R}$ it holds that

$$\alpha|_{\mathcal{O}(p)} = e^h \left(dt - \sum_{k=1}^{n-1} z_{n-1+k} dz_k - s_2 ds_1 \right), \quad (3.14)$$

and the Reeb field $R_\alpha = \frac{\partial}{\partial t}$;

- $W \cap \mathcal{O}(p) \ni q \mapsto (t(q), z_1(q), \dots, z_{2n-2}(q))$ is a local contact coordinate system around p in the relatively open neighborhood $W \cap \mathcal{O}(p)$ and

$$\alpha|_{W \cap \mathcal{O}(p)} = e^{h_0} \left(dt - \sum_{k=1}^{n-1} z_{n-1+k} dz_k \right), \quad (3.15)$$

where $h_0 = h|_{W \cap \mathcal{O}(p)}$. Moreover the Reeb field of $\alpha|_{W \cap \mathcal{O}(p)}$ is given by the restriction of $\frac{\partial}{\partial t}$ to $W \cap \mathcal{O}(p)$.

For convenience denote by $z_0 = t$. In the corresponding local coordinate vector fields

$$\frac{\partial}{\partial z_0} = \frac{\partial}{\partial t}, \frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2}, \dots, \frac{\partial}{\partial z_{2n-2}}, \frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_2}$$

we have

$$\hat{g} = \sum_{k,l=0}^{n-1} (g|_W)_{kl} dz_k \otimes dz_l + ds_1 \otimes ds_2. \quad (3.16)$$

It is easily computed that

$$\hat{g} \left(\nabla_{\frac{\partial}{\partial z_k}} \frac{\partial}{\partial z_l}, \frac{\partial}{\partial s_i} \right) = \frac{1}{2} (\hat{g}_{z_k s_i, z_l} + \hat{g}_{z_l s_i, z_k} - \hat{g}_{z_k z_l, s_i}) = 0.$$

So the second fundamental form of $W \cap \mathcal{V}$ with respect to $\hat{g}$ vanishes, that is, $W \cap \mathcal{V}$ is totally geodesics. $\square$

Proposition 3.3. *Let L be a compact Legendrian submanifold with boundary of the contact manifold (M, α) , and let W be a codimension two scaffold for L . Denote by $\hat{N}(L)$ the normal bundle of L with respect to $\hat{g}$. For $p \in \partial L$, suppose that $\hat{V} \in \hat{N}_p(L)$ satisfy the boundary condition*

$$(d\alpha)_p(N(p), \hat{V}) = 0.$$

Then $\hat{V} \in \xi'_p$. (In fact we have proved

$$\{\hat{V} \in \hat{N}_p(L) \mid (d\alpha)_p(N(p), \hat{V}) = 0\} \cap T_p\partial L = \{0\}.)$$

Proof. For any point $p \in \partial L$, take the local coordinate system around it as in the proof of Proposition 3.2. By composing with a suitable linear contactomorphism of form

$$\begin{aligned} \mathbb{R} \times \mathbb{R}^{2n-2} \times \mathbb{R}^2 &\rightarrow \mathbb{R} \times \mathbb{R}^{2n-2} \times \mathbb{R}^2 \\ (z_0, z_1, \dots, z_{2n-2}, s_1, s_2) &\mapsto (z_0, \mathcal{A}(z_1, \dots, z_{2n-2}), s_1, s_2), \end{aligned}$$

we may assume that

$$\left. \frac{\partial}{\partial z_1} \right|_p, \dots, \left. \frac{\partial}{\partial z_{n-1}} \right|_p$$

forms a basis of $T_p\partial L$, and that they are also orthogonal to vectors

$$\left. \frac{\partial}{\partial z_n} \right|_p, \dots, \left. \frac{\partial}{\partial z_{2n-2}} \right|_p$$

with respect to $\hat{g}$. (**Note:** Such a transformation does not change the Reeb field, i.e. we have still $R_\alpha = \frac{\partial}{\partial t}$.) Since the normal vector N of ∂L in L in the local coordinate system is equal to $\frac{\partial}{\partial s_1}$, we get an orthogonal basis of T_pL ,

$$\left. \frac{\partial}{\partial z_1} \right|_p, \dots, \left. \frac{\partial}{\partial z_{n-1}} \right|_p, \left. \frac{\partial}{\partial s_1} \right|_p.$$

It is easy to see that for some $\lambda \in \mathbb{R}$ the vector fields

$$\left. \frac{\partial}{\partial z_n} \right|_p, \dots, \left. \frac{\partial}{\partial z_{2n-2}} \right|_p, \left. \frac{\partial}{\partial s_2} \right|_p + \lambda \left. \frac{\partial}{\partial t} \right|_p$$

spans $\hat{N}_p(L)$, where $\hat{N}(L)$ the $\hat{g}$ -normal bundle of L . Let

$$\hat{V} := a_n \left. \frac{\partial}{\partial z_n} \right|_p + \dots + a_{2n-2} \left. \frac{\partial}{\partial z_{2n-2}} \right|_p + b \left(\left. \frac{\partial}{\partial s_2} \right|_p + \lambda \left. \frac{\partial}{\partial t} \right|_p \right) \in \hat{N}_p(L)$$

satisfy $(d\alpha)_p(N(p), \hat{V}) = 0$. Since $N(p) = \left. \frac{\partial}{\partial s_1} \right|_p$ and $R_\alpha(p) = \left. \frac{\partial}{\partial t} \right|_p$ we have

$$(d\alpha)_p \left(\left. \frac{\partial}{\partial s_1} \right|_p, a_n \left. \frac{\partial}{\partial z_n} \right|_p + \dots + a_{2n-2} \left. \frac{\partial}{\partial z_{2n-2}} \right|_p + b \left. \frac{\partial}{\partial s_2} \right|_p \right) = 0. \quad (3.17)$$

By (3.14) it is easy computed that

$$\begin{aligned} d\alpha|_{\mathcal{O}(p)} &= e^h \left(- \sum_{k=1}^{n-1} dz_{n-1+k} \wedge dz_k - ds_2 \wedge ds_1 \right) \\ &= e^h dh \wedge \left(dt - \sum_{k=1}^{n-1} z_{n-1+k} dz_k - s_2 ds_1 \right). \end{aligned} \quad (3.18)$$

Note that $s_1 = s_2 = 0$ at p . It follows from (3.17)-(3.18) that $be^h|_p = 0$ and thus $b = 0$. This shows

$$\hat{V} = a_n \frac{\partial}{\partial z_n} \Big|_p + \cdots + a_{2n-2} \frac{\partial}{\partial z_{2n-2}} \Big|_p \in \zeta'_p.$$

Clearly, when $\hat{V} \neq 0$ we have also $\hat{V} \notin T_p \partial L$. $\square$

Remark 3.4. Let $\widehat{\exp}$ be the exponent map of the metric $\hat{g}$. For any $p \in \partial L$ and $v \in \hat{N}_p(L)$ with $d\alpha(N(p), v) = 0$, Proposition 3.2 and Proposition 3.3 show that $\widehat{\exp}(p, v) \in W \cap \mathcal{U}$ if $|v|$ is small enough.

4. THE PROOF OF THEOREM 1.1

Let us start with a brief review of notations in Hodge theory (cf. [12, 15] for details) and then proceed to the proof of Theorem 1.1.

For $k \in \mathbb{N} \cup \{0\}$, $1 \leq p < \infty$ and $0 < a < 1$, let $W^{k,p}\Omega^r(L)$ (resp. $C^{k,a}\Omega^r(L)$) denote the space of r -forms of class $W^{k,p}$ (resp. $C^{k,a}$) as usual (cf. [12, 15]). Each form ω of them has a “tangential component” $\mathbf{t}\omega$ and a “normal component” $\mathbf{n}\omega$ (cf. [11, Def.4.2] or [15, (2.25)]), which satisfy

$$\mathbf{t}(\star\omega) = \star(\mathbf{n}\omega) \quad \text{and} \quad \mathbf{n}(\star\omega) = \star(\mathbf{t}\omega) \quad (4.1)$$

by Lemma 4.2 of [11], where $\star$ is the Hodge star operator of the metric $\hat{g}$. Set

$$\begin{aligned} C^{k,a}\Omega_{\mathbf{D}}^r(L) &:= \{\omega \in C^{k,a}\Omega^r(L) : \mathbf{t}\omega = 0\}, \\ C^{k,a}\Omega_{\mathbf{N}}^r(L) &:= \{\omega \in C^{k,a}\Omega^r(L) : \mathbf{n}\omega = 0\} \quad \text{and} \\ \mathcal{H}C^{k,a}\Omega^r(L) &:= \{\omega \in C^{k,a}\Omega^r(L) : d\omega = \delta\omega = 0\}. \end{aligned}$$

Replacing $C^{k,a}$ by $W^{k,p}$ gives corresponding spaces $W^{k,p}\Omega_{\mathbf{D}}^r(L)$, $W^{k,p}\Omega_{\mathbf{N}}^r(L)$ and $\mathcal{H}W^{k,p}\Omega^r(L)$. Clearly, for $S_{\mathbf{N}}^r = C^{k,\alpha}\Omega_{\mathbf{N}}^r(L)$ and $S_{\mathbf{D}}^r = C^{k,\alpha}\Omega_{\mathbf{D}}^r(L)$ (or $S_{\mathbf{N}}^r = W^{k,p}\Omega_{\mathbf{N}}^r(L)$ and $S_{\mathbf{D}}^r = W^{k,p}\Omega_{\mathbf{D}}^r(L)$), (4.1) implies

$$\star(S_{\mathbf{N}}^r) \subset S_{\mathbf{D}}^{n-r} \quad \text{and} \quad \star(S_{\mathbf{D}}^r) \subset S_{\mathbf{N}}^{n-r}. \quad (4.2)$$

By the definition of the co-differential δ , for any r -form ω it holds that

$$\star(\star\omega) = (-1)^{r(n-r)}\omega, \quad \star\delta\omega = (-1)^r d\star\omega, \quad \star d\omega = (-1)^{r+1} \delta\star\omega. \quad (4.3)$$

For $k \in \mathbb{N} \cup \{0\}$ the closure $C^{k,a}(d\Omega^r(L))$ of $d\Omega^r(L)$ in $C^{k,a}\Omega^{r+1}(L)$ is contained $\{d\eta : \eta \in C^{k+1,a}\Omega^r(L)\}$ by the Poincaré Lemma (cf. §3.1 of [1]).

Since the exponent map $\widehat{\exp}$ of the metric $\hat{g}$ is a local diffeomorphism, (by Tubular Neighborhood Theorem) the sufficiently small neighborhood of the zero section of $\hat{N}(L)$ satisfying the boundary condition corresponds to the deformations of submanifold L with boundary ∂L confined in W in one-to-one way.

Let $C^{2,a}(\Gamma(\widehat{N}(L)))$ denote the Banach space of $C^{2,a}$ -sections of the bundle $\widehat{N}(L)$. Define the Banach space

$$\mathcal{X} := \{V \in C^{2,a}(\Gamma(\widehat{N}(L))) : d\alpha(N, V|_{\partial L}) = 0\},$$

and denote by $\mathcal{U}$ a neighborhood of 0 in $\mathcal{X}$. Let $\widehat{\exp}$ be the exponent map of the metric $\hat{g}$. For $V \in \mathcal{U}$ define $\widehat{\exp}_V : L \rightarrow M$, $x \mapsto \widehat{\exp}_V(x) := \widehat{\exp}_x(V(x))$. Set

$$\begin{aligned} F : \mathcal{U} &\rightarrow C^{1,a}\Omega^1(L) \oplus C^{0,a}\Omega^n(L), \\ V &\mapsto ((\widehat{\exp}_V)^*\alpha, 2(\widehat{\exp}_V)^*\text{Im}\epsilon). \end{aligned} \quad (4.4)$$

It is C^1 as done in [1, 17]. Clearly, $\widehat{\exp}_V$ is homotopic to the inclusion $j : L \hookrightarrow M$ via $\widehat{\exp}_{tV}$, and hence they induce the same homomorphisms between the de Rham cohomology groups. It follows that the de Rham cohomology classes

$$[\widehat{\exp}_V^*(\text{Im}\epsilon)] = \widehat{\exp}_V^*[\text{Im}\epsilon] = j^*[\text{Im}\epsilon] = [j^*(\text{Im}\epsilon)] \in H^n(L, \mathbb{R}) \quad \text{vanish.}$$

This shows that

$$\text{Im}F \subseteq C^{1,a}\Omega^1(L) \oplus dC^{1,a}\Omega^{n-1}(L).$$

Consider F as a map to $C^{1,a}\Omega^1(L) \oplus dC^{1,a}\Omega^{n-1}(L)$.

To compute the differential of F at 0, for $V \in \mathcal{X}$ we set $f = \alpha(V)$ and $Y := V - fR_\alpha$. Then $f \in C^{2,a}(L)$ and $Y \in C^{1,a}(\Gamma(\xi|_L))$. By Proposition 3.3, $V(p) \in \xi'_p$ for any $p \in \partial L$, and so $f(p) = 0 \forall p \in \partial L$. Now $V = fR_\alpha + Y$. By the Cartan formula one can compute the linearization of F at 0,

$$\begin{aligned} F'(0)(V) &= \frac{d}{dt}(\widehat{\exp}_{tV}^*\alpha, 2\widehat{\exp}_{tV}^*\text{Im}\epsilon)|_{t=0} \\ &= (\mathcal{L}_V\alpha, 2\mathcal{L}_V\text{Im}\epsilon)|_L \\ &= (d\iota_{fR_\alpha+Y}\alpha + \iota_{fR_\alpha+Y}d\alpha, 2d\iota_{fR_\alpha+Y}\text{Im}\epsilon)|_L \\ &= (df + \iota_Y d\alpha, 2d\iota_Y \text{Im}\epsilon)|_L \\ &= (df + \iota_Y d\alpha, -d \star \iota_Y d\alpha)|_L \\ &= (d(f \circ j) + j^*(\iota_Y d\alpha), -d \star j^*(\iota_Y d\alpha)). \end{aligned} \quad (4.5)$$

Here the fifth equality comes from (2.2). In order to show that $F'(0)$ is surjective, we need to write each

$$(\eta, d\zeta) \in C^{1,\alpha}\Omega^1(L) \oplus dC^{1,\alpha}\Omega^{n-1}(L)$$

as a convenient form.

Note that $\mathbf{t}(d\omega) = d(\mathbf{t}\omega)$ and $\mathbf{n}(\delta\omega) = \delta(\mathbf{n}\omega)$ for any C^1 -form ω on L (cf. [15, Prop. 1.2.6(b)]). Since $C^{0,a}\Omega^{n-1}(L) \subset L^2\Omega^{n-1}(L)$, by [11, Th.5.7, 5.8] or [12, Th.7.7.7, 7.7.8] we may write $\zeta = \delta_n\gamma' + d\gamma'' + h(\zeta)$, where

$$\gamma' \in C^{1,a}\Omega_{\mathbf{N}}^n(L), \quad \gamma'' \in C^{1,a}\Omega_{\mathbf{D}}^{n-2}(L), \quad h(\zeta) \in \mathcal{H}C^{0,a}\Omega^{n-1}(L).$$

Moreover (4.2) and $\mathbf{t}(d\omega) = d(\mathbf{t}\omega)$ imply

$$\delta_n = (-1)^{n(n+1)+1} \star d_0 \star : C^{2,\alpha} \Omega_{\mathbf{N}}^n(L) \rightarrow C^{1,\alpha} \Omega_{\mathbf{N}}^{n-1}(L).$$

We may assume

$$d\zeta = -d \star du \quad \text{with} \quad u \in C^{2,\alpha} \Omega_{\mathbf{D}}^0(L).$$

Similarly, we have

$$\eta = \delta v + d\beta + h(\eta),$$

where

$$v \in C^{2,\alpha} \Omega_{\mathbf{N}}^2(L), \quad \beta \in C^{2,\alpha} \Omega_{\mathbf{D}}^0(L), \quad h(\eta) \in \mathcal{H} C^{1,\alpha} \Omega^1(L).$$

By (4.3), $d \star \delta v = (-1)^2 d(d \star v) = 0$ and $d(\star h(\eta)) = (-1) \star \delta h(\eta) = 0$. We get

$$\begin{aligned} (\eta, d\zeta) &= (d\beta - du + du + \delta v + h(\eta), -d \star (du + \delta v + h(\eta))) \\ &= (d\chi + \omega, -d \star \omega), \end{aligned}$$

where

$$\begin{aligned} \chi &:= \beta - u \in C^{2,\alpha} \Omega_{\mathbf{D}}^0(L), \\ \omega &:= du + \delta v + h(\eta) \in C^{1,\alpha} \Omega^1(L). \end{aligned} \tag{4.6}$$

Take $f = \chi$. We need to find a $Y \in C^{1,a}(\Gamma(\xi|_L))$ such that

$$f R_\alpha + Y \in C^{1,a}(\Gamma(\widehat{N}(L))) \quad \text{and} \quad j^*(\iota_Y d\alpha) = \omega.$$

Since $j^*(\iota_{Y+fR_\alpha} d\alpha) = j^*(\iota_Y d\alpha) = \omega$, it suffices to find a $Z \in C^{1,a}(\Gamma(\widehat{N}(L)))$ such that

$$j^*(\iota_Z d\alpha) = \omega. \tag{4.7}$$

To this goal, consider the symplectic vector bundle $(\xi|_L, d\alpha|_{\xi|_L})$ with a Lagrangian subbundle TL . Let $TL_\xi^{\perp \hat{g}}$ be the orthogonal complementary bundle of TL in $\xi|_L$ with respect to $\hat{g}$. Then $TL_\xi^{\perp \hat{g}} = \xi \cap \widehat{N}(L)$. So $\xi|_L = TL \oplus_{\hat{g}} (\xi \cap \widehat{N}(L))$. Note that ω may be viewed as a section of the bundle $\text{Hom}(TL, \mathbb{R})$. We may extend it into a section of $\text{Hom}(\xi|_L, \mathbb{R})$, $\hat{\omega}$, by defining

$$\hat{\omega}_p(u+v) = \omega_p(u)$$

for any $p \in L$ and $u+v \in T_p L \oplus_{\hat{g}} (\xi \cap \widehat{N}(L))_p$, where $u \in T_p L$ and $v \in (\xi \cap \widehat{N}(L))_p$. Note that $\hat{\omega} \in C^{1,a}(\Gamma(\text{Hom}(\xi|_L, \mathbb{R})))$. The non-degeneracy of $d\alpha$ on ξ implies that there exists a unique section $Z : L \rightarrow \xi|_L$ such that

$$(d\alpha)_p(Z(p), A) = \hat{\omega}_p(A) \quad \forall p \in L \text{ and } A \in \xi_p.$$

Clearly, $Z \in C^{1,a}(\xi|_L)$. Since $\xi|_L = TL \oplus_{\hat{g}} (\xi \cap \widehat{N}(L))$ we get a unique decomposition $Z = Z_1 + Z_2$, where $Z_1 \in C^{1,a}(\Gamma(TL))$ and $Z_2 \in C^{1,a}(\Gamma(\xi \cap \widehat{N}(L)))$. Obverse that

$$j^*(\iota_{Z_1} d\alpha) = 0.$$

In fact, for any $p \in L$ and $u \in T_p L$ it holds that

$$(j^*(\iota_{Z_1} d\alpha))_p(u) = (\iota_{Z_1} d\alpha)_{j(p)}(j_* u) = (d\alpha)_p(Z_1(p), u) = 0$$

since $T_p L$ is a Lagrangian subspace of $(\xi_p, (d\alpha)_p)$. Hence we get

$$(d\alpha)_p(Z_2(p), A) = \hat{\omega}_p(A) \quad \forall p \in L \text{ and } A \in \xi_p.$$

This implies $j^*(\iota_{Z_2} d\alpha) = \omega$. In summary we have proved:

Claim 4.1. *There exists a unique section $Z : L \rightarrow \xi|_L \cap \hat{N}(L)$ such that (4.7) is satisfied. Moreover, Z is also of class $C^{1,a}$. As a consequence the map $F'(0)$ is surjective.*

Next let us compute $\ker(F'(0))$. Let $V \in \mathcal{X}$ sit in $\ker(F'(0))$. As above we may write $V = fR_\alpha + Y$, where $f = \alpha(V)$ and $Y \in C^{k+1,a}(\Gamma(\xi|_L))$. (4.5) yields

$$df + j^*(\iota_Y d\alpha) = 0, \quad (4.8)$$

$$-d \star j^*(\iota_Y d\alpha) = 0. \quad (4.9)$$

From (4.8) we get

$$\begin{aligned} 0 = \delta(df + j^*(\iota_Y d\alpha)) &= \delta df + \delta(j^*(\iota_Y d\alpha)) \\ &= \delta df + (-1)^{2n+1} \star d \star (j^*(\iota_Y d\alpha)) \\ &= \delta df \end{aligned}$$

because of (4.9). Hence $\Delta f = 0$. By Proposition 3.3 the boundary condition $d\alpha(N, V|_{\partial L}) = 0$ implies $V(p) \in \xi'_p \subset \xi_p$ for any $p \in \partial L$, and thus $f|_{\partial L} = 0$. Since ∂L is a non-empty closed manifold, the maximum principle leads to $f \equiv 0$. Hence $V = Y$. By (4.8) we have

$$(d\alpha)_p(Y(p), u) = 0 \quad \forall p \in L \text{ and } u \in T_p L.$$

This means that $Y(p)$ belongs to $T_p L$ since $T_p L$ is a Lagrangian subspace in $(\xi_p, (d\alpha)_p)$. Moreover, $Y(p) = V(p) \in \hat{N}_p(L)$. Hence we get $V(p) = 0$ for any $p \in L$. It shows

$$\ker F'(0) = 0.$$

Combing this with Claim 4.1 we prove that the differential

$$F'(0) : T_0 \mathcal{X} \rightarrow C^{1,a}\Omega^1(L) \oplus dC^{1,a}\Omega^{n-1}(L)$$

is a Banach space isomorphism. The inverse function theorem implies that there exists a neighborhood of 0 in $\mathcal{X}$, $\mathcal{U}_0 \subset \mathcal{U}$, such that $F^{-1}(0) \cap \mathcal{U}_0 = \{0\}$. This completes the proof of Theorem 1.1. $\square$

5. THE PROOF OF THEOREM 1.2

Let $\langle R_\alpha \rangle$ denote the real line bundle generated by $R_\alpha|_L$. Then the normal bundle of L with respect to the metric g , $N(L)$, is equal to $\langle R_\alpha \rangle \oplus_g JTL$. For a small section $V : L \rightarrow \mathcal{N}(L)$, the exponent map of g yields a map

$$\exp_V : L \rightarrow M, \quad x \mapsto \exp_x(V(x)).$$

Thus there exists a neighborhood $\mathcal{V}$ of 0 in

$$\mathcal{Y} := \{V \in C^{2,a}(\Gamma(\langle R_\alpha \rangle)) \oplus C^{1,a}(\Gamma(JTL)) : \alpha(V)|_{\partial L} = \text{const}\}$$

so that the following map is well-defined:

$$\begin{aligned} G : \mathcal{V} &\rightarrow C^{1,a}\Omega^1(L) \oplus C^{0,a}\Omega^n(L), \\ V &\mapsto (\exp_V^* \alpha, 2 \exp_V^* \text{Im} \epsilon). \end{aligned} \quad (5.1)$$

It is C^1 ([17]), and $\text{Im}(G) \subseteq C^{1,a}\Omega^1(L) \oplus dC^{1,a}\Omega^{n-1}(L)$ as above since $\exp_V$ is homotopic to the inclusion $j : L \hookrightarrow M$ via $\exp_{tV}$.

Considering G as a map to $C^{1,a}\Omega^1(L) \oplus dC^{1,a}\Omega^{n-1}(L)$, and writing $V = JX + fR_\alpha$, we may get

$$\begin{aligned} G'(0)(V) &= \frac{d}{dt}(\exp_{tV}^* \alpha, 2 \exp_{tV}^* \text{Im} \epsilon)|_{t=0} \\ &= (\mathcal{L}_V \alpha, 2\mathcal{L}_V \text{Im} \epsilon)|_L \\ &= (df + \iota_{JX} d\alpha, -d * \iota_{JX} d\alpha)|_L \end{aligned} \quad (5.2)$$

as above. Moreover, each $(\eta, d\zeta) \in C^{1,a}\Omega^1(L) \oplus dC^{1,a}\Omega^{n-1}(L)$ may be written as $(\eta, d\zeta) = (d\chi + \omega, -d\star\omega)$, where χ and ω are as in (4.6). Take $f = \chi$, and one easily find $X \in C^{1,\alpha}(\Gamma(TL))$ such that $j^*(\iota_{JX} d\alpha) = \omega$. Clearly, such a $V = fR_\alpha + JX$ satisfies $\alpha(V)|_{\partial L} = 0$. Hence $G'(0)$ is surjective.

Assume that $V = fR_\alpha + JX$ sits in $\ker(G'(0))$. Then f and JX satisfy

$$df + j^*(\iota_{JX} d\alpha) = 0, \quad -d\star j^*(\iota_{JX} d\alpha) = 0.$$

It follows that $\triangle f = \delta df = 0$. Recall that $f = \alpha(V)$ is equal to a constant c on ∂L . By the maximum principle we get $f \equiv c$, and hence

$$j^*(\iota_{JX} d\alpha) = 0.$$

From this we derive $JX = 0$ as above. This prove $\ker(G'(0)) = \{cR_\alpha \mid c \in \mathbb{R}\}$. Hence $(0, 0)$ is a regular value of the restriction of G to a small neighborhood of $0 \in \mathcal{V}$, and thus the moduli space $\mathfrak{M}(L)$ is a 1-dimensional smooth manifold by the implicit function theorem. $\square$

Remark 5.1. If we replace $\mathcal{V}$ by a neighborhood $\mathcal{W}$ of 0 in

$$C^{2,a}(\Gamma(\langle R_\alpha \rangle)) \oplus C^{1,a}(\Gamma(JTL)),$$

then the map

$$\widehat{G} : \mathcal{W} \rightarrow C^{1,a}(\Lambda^1(L)) \oplus C^{0,a}(\Lambda^n(L)), \quad V \mapsto (\exp_V^* \alpha, 2 \exp_V^* \operatorname{Im} \epsilon).$$

is still C^1 and has the image $\operatorname{Im}(\widehat{G}) \subseteq C^{1,a}(\Lambda^1(L)) \oplus dC^{1,a}(\Lambda^{n-1}(L))$. From the above proof it is easy to see that $\widehat{G}'(0)$ is surjective. If $V = fR_\alpha + JX$ belongs to $\ker(\widehat{G}'(0))$, we have $\triangle f = 0$ as above. But ∂L is a nonempty closed manifold, by Theorem 3.4.6 of [15] each $b \in C^\infty(\partial L)$ corresponds to a unique $f \in C^\infty(L)$ satisfying $\triangle f = 0$ and $f|_{\partial L} = b$. It follows that $\ker(\widehat{G}'(0))$ must be infinitely dimensional.

Starting from a closed special Legendrian submanifold one may construct a lot of compact special Legendrian submanifolds satisfying our theorems above by Darboux theorem in contact geometry. Moreover, the corresponding problems with [1, Cor.9] and [17, Th.4.8] can also be considered similarly.

REFERENCES

- [1] A. Butsher, Deformations of minimal Lagrangian submanifolds with boundary, *Proc. Amer. Math. Soc.* **131**(6) (2002) 1953-1964.
- [2] P. De Bartolomeis, Geometric Structures on Moduli Spaces of Special Lagrangian Submanifolds, *Ann. di Mat. Pura ed Applicata*, IV, Vol. CLXXIX (2001), pp. 361–382.
- [3] D. Gayet and F. Witt, Deformations of associative submanifolds with boundary, *Adv. in Math.*, **226**(2011), 2351-2370.
- [4] H. Geiges, An Introduction to Contact Topology Cambridge: Cambridge University Press, 2008.
- [5] F.R. Harvey and H. B. Lawson, Calibrated Geometries, *Acta. Math.*, **148**(1982)47-157.
- [6] D.D. Joyce, Riemannian Holonomy Group and Calibrated Geometry. Oxford: Oxford University Press, 2007.
- [7] D.D. Joyce, Special Lagrangian Submanifolds with Isolated Conical Singularities. II. Moduli Spaces, *Annals of Global Analysis and Geometry*, **25**(2004), 301C352.
- [8] A. Kovalev and J. D. Lotay, Deformations of compact coassociative 4-folds with boundary, *J. Geom. phy.* **59** (2009)63-73.

- [9] D. McDuff and D. Salamon, Introduction to Symplectic Topology (second edition). Oxford: Clarendon Press, 1998.
- [10] R. C. McLean, Deformation of calibrated submanifolds. *Comm Anal Geom.* **6**(1998), 705-747.
- [11] C. B. Morrey, A variational method in the theory of harmonic integrals II. *Amer. J. Math.* **78**(1956), 137-170.
- [12] C. B. Morrey, Multiple Integrals in the Calculus of Variations. Grundlehren der mathematischen Wissenschaften 130, Springer Verlag, Berlin 1966.
- [13] T. Pacini, Special Lagrangian conifolds, I: moduli spaces, arXiv:1002.1222v1 [math.DG].
- [14] S. Salur, Deformation of special Lagrangian submanifolds. *Communication in Contemporary Mathematics.* **2**(2000), no 3, 365-372.
- [15] G. Schwarz, Hodge Decomposition—A Method for Solving Boundary Value Problems, *Springer-Verlag*, Berlin, 1995. MR **96k**:58222.
- [16] J. Simons, Minimal varieties in Riemannian manifolds, *Ann. of Math.*, **88**(1968), 62-105.
- [17] A. Tomassini and L. Vezzoni, Contact Calabi-Yau manifolds and special Legendrian submanifolds, *Osaka J. Math.*, **45**(2008), 127-147.

SCHOOL OF MATHEMATICAL SCIENCES, BEIJING NORMAL UNIVERSITY, LABORATORY OF MATHEMATICS AND COMPLEX SYSTEMS, MINISTRY OF EDUCATION, BEIJING 100875, THE PEOPLE'S REPUBLIC OF CHINA.

E-mail address: gclu@bnu.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, BEIJING NORMAL UNIVERSITY, BEIJING 100875, THE PEOPLE'S REPUBLIC OF CHINA.

E-mail address: xmchen1983@mail.bnu.edu.cn